


















WILDLIFE ON AN ENHANCED FARM DAM IN SOUTH-EASTERN AUSTRALIA








SUSTAINABLE
FARMS





-  Microbat, southern myotis, *Myotis macropus*
-  Common brown butterfly, *Heteronympha merope*
-  Golden-headed cisticola, *Cisticola exilis*
-  Purple swamphen, *Porphyrio porphyrio*
-  Buff-banded rail, *Gallirallus philippensis*

-  Australian reed warbler, *Acrocephalus australis*
-  Blue skimmer dragonfly, *Orthetrum caledonicum*
-  Black-winged stilt, *Himantopus himantopus*
-  Peron's tree frog, *Litoria peronii*
-  Sudell's frog, *Neobatrachus sudellae*

-  Superb fairy wren, *Malurus cyaneus*
-  Australian wood ducks, *Chenonetta jubata*
-  Latham's snipe, *Gallinago hardwickii*
-  Australasian grebe, *Tachybaptus novaehollandiae*
-  Eastern long-necked turtle, *Chelodina longicollis*

-  Eastern great egret, *Ardea modesta*
-  Welcome swallow, *Hirundo neoxena*
-  Australian shelducks, *Tadorna tadornoides*
-  Spotted marsh frog, *Limodynastes tasmaniensis*
-  Freshwater yabby, *Cherax destructor*

-  Swamp harrier, *Circus approximans*
-  Diamond firetail, *Stagonopleura guttata*
-  White-faced heron, *Egretta novaehollandiae*
-  Yellow-billed spoonbill, *Platalea flavipes*
-  Black-fronted dotterel, *Eileynis melanops*

-  Red-browed finch, *Neochmia temporalis*
-  Fairy martin, *Petrochelidon ariel*
-  White-necked heron, *Ardea pacifica*
-  Rakali, *Hydromys chrysogaster*